TRANSACTIONS

OF THE

BISHOP'S STORTFORD & DISTRICT NATURAL HISTORY SOCIETY

VOL. I

LIST OF

VASCULAR PLANTS

OF

BISHOP'S STORTFORD AND DISTRICT

COMPILED BY MEMBERS OF THE SOCIETY

1952

PUBLISHED FOR THE SOCIETY PRICE 58.

TRANSACTIONS

OF THE

BISHOP'S STORTFORD AND DISTRICT NATURAL HISTORY SOCIETY

Vol. I Part II

THE VASCULAR PLANTS OF
BISHOP'S STORTFORD AND DISTRICT

FOREWORD

WHEN our Society was founded in 1935 it was decided that our "district" should comprise the area included by a circle having a radius of six miles centred on the Corn Exchange, Bishop's Stortford, This area includes the following villages and hamlets:

Albury	Hatfield Broad Oak	Quendon
Berden	Hatfield Heath	Rickling
Birchanger	Henham	Sawbridgeworth
Braughing	High Wych	Sheering
Elsenham	Little Hadham	Standon
Farnham	Little Hallingbury	Stocking Pelham
Furneux Pelham	Manuden	Takeley
Gilston	Much Hadham	Ugley
Great Hallingbury	Patmore Heath	Widford .

Over much of this area the soil is boulder clay; this is the home of the Oxlip, *Primula elatior*, and in many of our woods it occurs in great abundance and provides a most beautiful sight in April. Where the chalk nears the surface or crops out, typically lime-loving plants are found in limited numbers. The dominant trees are oak and hornbeam.

Up to the present (1950) some parts of this area have not been thoroughly hunted and there are gaps in our knowledge of some Orders, especially *Cyperaceae*. The list must therefore be regarded as only a tentative one. A number of species not included are known to occur just outside our district. When the whole area has been thoroughly worked it is likely that a number of additions will have to be made.

The list has been compiled by the following members of our Society to which their ciphers are appended:

B. J. Adams (As)	H. J. Holmes (Hs)
P. B. M. Allan (An)	Mrs. Kedge (Kg)
R. D. Anstead (Ad)	J. Killick (Kk)
C. S. Colman (Cn)	N. Monk-Jones (MJ)
C. Craufurd (Cd)	Miss L. W. Laverick (Lk)
E. G. Eagling (Eg)	W. E. Palmer (Pr)
Miss M. Elliot (Et)	Miss Rawlins (Rs)
Mrs. N. Ferguson (Fn)	Miss Turner (Tr)
A. E. Hick (Hk)	

The discovery in our district of species not in this list should be notified with specimens to Mr. R. D. Anstead, 5 Maple Avenue, Bishop's Stortford.

Page Three

The Cheshunt Press Ltd., Cheshunt, and Embassy Buildings, Waltham Cross.

The nomenclature adopted in this list is that in the Check-List of British Vascular Plants compiled by Prof. A. R. Clapham and issued by the British Ecological Society in 1946 (Journal of Ecology, 333, pp. 308 seq.).

Where the botanical name of a plant is followed by "agg." it indicates that the species has now been broken up by some authorities into a number of sub-species and varieties whose claim to specific rank is not always generally conceded. A more critical determination of these has not yet been possible in our area, and more intensive study of them is required.

31st December, 1950.

Ranunculaceae

- I. Clematis vitalba L. : TRAVELLER'S JOY
 Ubiquitous throughout our district, in hedgerows and woods.
 Occasionally attains a large size with thick "liana-like" stems.
- 2. Thalictrum flavum L.: Yellow Meadow Rue First found 10.vii.20 in meadows by Rye Street and specimens brought from there since (Cn). Also in meads by the River Stort.
- 3. Anemone nemorosa L.: Wood Anemone
 Very common in all woods in our district, spreading into grassland on the verges. Whenever woods are coppiced and the light let in
 the anemones spread very rapidly and soon make a complete ground
 cover.
 - 4. Adonis annua L. : PHEASANT'S EYE Found once only, 1938 (Kg).
- 5. Ranunculus fluitans agg.: WATER CROWFOOT Common in ponds and streams. This plant has many forms which have been distinguished as species. The local forms need further study.
- 6. Ranunculus flammula L.: Lesser Spearwort
 Not very common. Woodside Green (Cn); Hatfield Forest; by
 the Stort and canal, and occasionally in other wet places.
- 7. Ranunculus ficaria L. : LESSER CELANDINE (PILEWORT)
 One of our commonest Spring flowers, occurring throughout the district.
- 8. Ranunculus sceleratus L.: CELERY-LEAVED CROWFOOT Frequent in our district in wet places. Bury Green in quantity, 1939-40 (Ad); Hatfield Forest, in the swamp.
- Ranunculus auricomus L.: Wood Crowfoot (Goldilocks)
 Frequent in our district in woods and shady hedgerows. Usually
 with deformed petals, but with complete corolla—Bury Green Lane,
 1939 (Ad).
 - 10. Ranunculus acris L. : MEADOW BUTTERCUP Plentiful everywhere in grasslands and by roadsides.
- Common in our district, often a troublesome garden weed. Favours damp places. The glades in Stocking Plantation are carpeted with it.

Page Five

- 12. Ranunculus bulbosus L. : Bulbous Buttercup Plentiful in grasslands. In some fields in this district it makes very striking carpets of bloom in early Spring.
- 13. Ranunculus parviflorus L.: SMALL-FLOWERED CROWFOOT London Bridge in Hatfield Forest, 1936. It has now (1949) spread over a large area there beside Elgin Coppice.
 - 14. Ranunculus arvensis L.: CORN CROWFOOT Common in cornfields throughout our district.
- 15. Caltha palustris L.: MARSH MARIGOLD (KINGCUPS)
 Occasional and widespread in wet places, but not in large quantities. Birchanger; by the Stort and canal; Hatfield Forest.
- 16. Helleborus viridis L.: GREEN HELLEBORE "Specimens brought to me from Takeley Woods by L. J. Bacon, 11.v.29" (Cn). Frequently grown in gardens.
- 17. Aconitum anglicum Stapf: Monkshood A long-established clump of this plant on the river bank above Pinchpools, Manuden, may be considered wild (An).

Berberidaceae

18. Berberis vulgaris L.: Common Barberry
Only one shrub of this is known in our district, in a hedge between Parsonage Lane and Birchanger. It will probably be destroyed
as the site is now a building area.

Nymphaeaceae

19. Nuphar lutea (L.) Sm. : Yellow WATERLILY In the Stort and canal and their backwaters.

Papaveraceae

- 20. Papaver rhoeas L. : COMMON RED POPPY Very frequent in cornfields and by roadsides.
- 21. Papaver dubium L. : Long Smooth-Headed Poppy Scarce. Waste ground off Maple Avenue, 1940 (Cn).
- 22. Chelidonium majus L. : CELANDINE Foxdells, 1938; Quendon, 1938; Manuden, 1938; Wallbury Dells, 1942.

Page Six

Fumariaceae

- 23. Corydalis lutea (L.) DC. : YELLOW CORYDALIS
 Quite frequent but probably always a garden escape and not truly
 wild. It is very commonly grown in cottage gardens throughout our
 district.
- 24. Fumaria officinalis L.: COMMON FUMITORY
 Frequent in our district. This plant needs more careful study as
 there are quite possibly other species to be found.

Cruciferae

- 25. Nasturtium officinale R.Br.: WATERCRESS Common throughout the district in ditches and streams.
- 26. Nasturtium sylvestris (L.) Besser : Creeping Yellow Cress In Meadowdyke garden, Maze Green Road, as a persistent weed (Cn). Found nowhere else.
- 27. Nasturtium islandica (Oeder) Borbás : MARSH WATERCRESS
 By the canal near London Bridge, Bishop's Stortford, 1940; Sawbridgeworth, 1938; Rickling, 1938.
- 28. Nasturtium amphibia (L.) Besser: Great Watercress
 On the lock gates at Spellbrook, 1938 (Ad); on waste ground by
 London Bridge, Bishop's Stortford, 1941 (Ad).
- 29. Barbarea vuglaris R.Br. : COMMON WINTER CRESS (YELLOW ROCKET)

 Common throughout our district. Occurs in Hatfield Forest.
- 30. Cardamine pratensis L.: LADY'S SMOCK (CUCKOO FLOWER)
 Plentiful everywhere. A double-flowered variety was found at
 Hatfield Forest in 1947.
 - 31. Cardamine hirsuta L. : HAIRY BITTERCRESS A common weed in gardens and cultivated areas.
- 32. Erophila verna (L.) Chevall. : WHITLOW GRASS
 Rare in our district and reported from one locality only: Hatfield
 Forest, 1936 (Cn), 1941 (Ad).
- 33. Armoracia lapathifolia Gilib. : Horseradish Frequent by roadsides and in waste places. Probably usually a garden escape.

Page Seven

- 34. Arabidopsis thaliana (L.) Heynh.: Thalecress (Wallcress)
 Occasional. Twyford Road, 1938 (As); railway-bank near footbridge, Bishop's Stortford, 1941 (Ad); level-crossing, Birchanger footpath, 1944 (Ad).
 - 35. Sisymbrium officinale (L.) Scop. : HEDGE MUSTARD Very common throughout our district.
- 36. Descurainia sophia (L.) Prantl: FLIXWEED Found in one place only, the waste ground between the gasworks and the canal (Cn); 1940 (Ad); 1941 (Ad), (Cn).
- Alliaria petiolata (Bieb.) Cavara & Grande: GARLIC MUSTARD 37. Alliaria petiolata (Dico.) Carata Ubiquitous throughout our district.
- 38. Erysimum cheiranthoides L. : TREACLE MUSTARD Found in one locality only: Wickham Hall farm near the Farnham road. First reported in 1944 and seen each year since then.
 - 39. Brassica oleracea L. : WILD CABBAGE Common throughout our district, being often cultivated.
 - Brassica campestris L. : FIELD CABBAGE Stacey's Farm, 1939; field by Hadham road rifle butts, 1940 (Ad).
- 41. Brassica nigra (L.) Koch: COMMON MUSTARD In rough ground near the town; but birds probably carry the seed, as this plant is commonly cultivated as a green dressing crop.
 - Capsella bursa-pastoris (L.) Medic. : Shepherd's Purse One of the commonest weeds throughout the district.
- 43. Coronopus squamatus (Forsk.) Aschers. : WARTCRESS (SWINE'S CRESS)

 An uncommon plant in our district, though it may possibly be frequently overlooked. Manuden, 1938 (Ad); the College upper field,

44. Lepidium sativum L.: CRESS
This plant appeared as a weed in Mr. Sutton's garden, Maze
Green road, in July 1942.

Page Eight

- 45. Cardaria draba (L.) Desv.: HOARY CRESS
 This plant appears to be spreading and becoming quite common.
 Twyford level-crossing, 1938 (As); Maple Avenue, 1939 (Ad); waste ground by the river, 1941 (Ad); Cricketfield Lane, 1944 (Hk).
- 46. Thlaspi arvense L.: Penny-Cress (Mithridate Mustard) First recorded 16.vi.21 in the College grounds (Cn). Seen again in the same place, 1940-1947 (Cn). By the wall of the gasworks, on the canal towing-path, 1941 (Ad), (Cn).
 - 47. Raphanus raphanistrum L.: WILD RADISH A common weed of cultivation throughout our district.

Resedaceae

48. Reseda luteola L. : DYER'S ROCKET (YELLOW WEED, WELD)
Occasional; usually on chalky soil. Hatfield Forest, 1939* (Ad);
Wallbury Dells, 1939; Manuden, in a gravel pit, 1948 (Hk).

Cistaceae

49. Helianthemum nummularium (L.) Mill. : COMMON ROCKROSE
This plant occurs occasionally in our district where the chalk
actually appears on the surface and is strictly confined to such places.
By the footpath by Cherry Cottages near the Farnham road; Little
Hadham Farm, 1941 (Ad); near Broome Wood, 1948 (Et).

Violaceae

- 50. Viola odorata L. : SWEET VIOLET
 The blue-flowered sweet violet is quite rare in our district, but the white-flowered variety is very common everywhere in hedgerows.
- 51. Viola hirta L.: HAIRY VIOLET
 Only once recorded: Hatfield Forest (MJ). This plant probably does occur, however, in our district but escapes notice. A more diligent search for it is required.
- Very frequent throughout the district. Particularly fine displays of it are to be seen in woods such as Canfield Hart. It varies very much in the colour and markings of the flowers and has been divided by some authorities into a number of different species. For lack of study of these variations found locally all are included under the above

Page Nine

53. Viola tricolor agg. : HEARTSEASE
A common cornfield weed throughout our district. This again has several different varieties some of which have been given specific rank.

Polygalaceae

54. Polygala vulgaris L.: COMMON MILKWORT
Quite frequent in our district wherever the chalk nears the surface. Both white and blue varieties are commonly found and occasionally the red form. Hatfield Forest; Farnham gravel pit; Tom Wood's Pasture, and elsewhere.

Caryophyllaceae

- 55. Dianthus armeria L. : DEPTFORD PINK In one locality only: Bishop's Stortford golf links, 1942 (Hs), and each year since then.
 - 56. Silene cucubalus Wibel : BLADDER CAMPION Common throughout our district.
- 57. Silene gallica agg. : ENGLISH CATCHFLY Rare. Reported only once, from the College playing-fields, 1941 (As).
- 58. Melandrium noctiflorum (L.) Fr.: NIGHT-FLOWERING
 CATCHFLY
 Occasional. Piggot's Farm in a cornfield, 1940 (As); plentiful at
 Thorley, 1941 (As).
- 59. Melandrium album (Mill.) Garcke: WHITE CAMPION
 Very common throughout our district, especially in cornfields.
 Increasing in frequency as the soil becomes chalky on the Cambridge side.
 - 60. Melandrium dioicum (L.) Coss. & Germ. : RED CAMPION Ubiquitous throughout our district.
- 61. Lychnis flos-cuculi L.: RAGGED ROBIN
 Frequent in wet places. Brewery Fields; Broome Wood (An);
 Patmore Wood, Albury (An); Thorley Wash; Birchanger.

Page Ten

- 62. Agrostemma githago L.: CORN COCKLE

 Now very rare and seldom seen in our district from which it
 seems to have disappeared. The last time it was recorded was in 1938

 (Cd).
 - 63. Cerastium vulgatum L. : MOUSE-EAR CHICKWEED A ubiquitous weed.
- 64. Cerastium arvense L.: FIELD CHICKWEED

 Manuden; Rickling, 1938; in fields by the canal, etc.; Patmore
 Heath, 1950.
- 65. Myosoton aquaticum (L.) Moench: WATER STITCHWORT Frequent by the canal and its backwaters and apparently spreading.
 - 66. Stellaria media (L.) Vill. : CHICKWEED Plentiful everywhere.
 - 67. Stellaria holostea L. : Greater Stitchwort Plentiful everywhere.
 - 68. Stellaria graminea L. : LESSER STITCHWORT Very common throughout the district.
 - 69. Arenaria verna L.: VERNAL SANDWORT Not common. Reported from Hatfield Forest, 1939 (Tr).
 - 70. Arenaria trinervia L. : THREE-NERVED SANDWORT A very common plant in our district in hedgerows and woods.
- 71. Arenaria serpyllifolia L.: THYME-LEAVED SANDWORT On a wall at Manuden (Cn); Farnham gravel-pit at the base of a chalk cliff, 1944 (Ad); garden paths, Maple Avenue (Ad).
- 72. Sagina procumbens L.: PROCUMBENT PEARLWORT Common everywhere in waste and stony places. Often found on garden paths and rockeries.
- 73. Spergularia rubra (L.) J. & C. Presl : Purple Sandwort Rare. Seen once or twice on gravel paths in the College grounds (Cn).

Page Eleven

Portulacaceae

Claytonia perfoliata Willd. : PERFOLIATE CLAYTONIA 74. Claytonia perfotata Willd. : PERFOLIATE CLATTON BOODSTRUCTURE TO STATE OF THE PROPERTY OF Hadham Road in 1940-41 (Fn); also in a garden at Meadowdyke, Maze Green Road, some years ago (Cn).

75. Montia fontana L. : WATER BLINKS Patmore Heath, 1950 (Ad).

Tamaricaceae

76. Tamarix gallica L. : TAMARISK In gardens only; not wild in our district.

Hypericaceae

- Hypericum perforatum L. : PERFORATED ST. JOHN'S WORT Plentiful everywhere.
- 78. Hypericum quadrangulum L.: SQUARE-STALKED ST. JOHN'S WORT Frequent in woods in our district. Stocking Plantation; Thorley Wood; Hatfield Forest, etc.
 - 79. Hypericum pulchrum L. : Slender St. John's Wort Common everywhere in the district.
- 80. Hypericum hirsutum L. : HAIRY ST. JOHN'S WORT Frequent in woods in our district. Thorley Wood; Canfield Hart; Hatfield Forest, etc.

Malvaceae

- 81. Malva moschata L.: MUSK MALLOW Plentiful, especially on chalky soil. A white-flowered variety is sometimes found: Farnham chalk-pit, 1939-49.
 - 82. Malva sylvestris L. : Common Mallow A common roadside plant throughout the district.
 - 83. Malva rotundifolia L. : DWARF MALLOW Occasional but probably overlooked.

Page Twelve

84. Abutilon indicum. G. Don.
An introduced plant which appeared in a garden in Stansted Road in 1942 (Hk). On an allotment in Rye Street in 1943 (Cd). On an allotment near the Red White and Blue inn in 1945 and 1949 (Cd).

85. Hibiscus trionum L.
A weed of cultivated and waste places, very common in cornfields in E. Europe and thus imported into Britain with corn, appearing appradically where corn siftings are thrown. Wickham Hall Farm in a field of lucerne, 1949 (Cd).

Tiliaceae

86. Tilia vulgaris Hayne : LIME Not wild, but many avenues have been planted with it in the town.

Linaceae

- 87. Linum catharticum L. : CATHARTIC FLAX Very common on chalky soil in our district. Farnham gravel-pit; Randall's Farm, Takeley; Hatfield Forest, etc.
- 88. Linum bienne Mill. : NARROW-LEAVED FLAX Upper Field, the College grounds, 1909 (Cn). Not reported since then.

Geraniaceae

89. Geranium sanguineum L. : BLOODY CRANE'S-BILL Found in our district once only, at Pedler's Green (MJ).

90. Geranium versicolor L. A garden escape; found in two places at Hockerill in 1946.

- 91. Geranium phaeum L. : DUSKY CRANE'S-BILL Doubtfully wild in our district; reported once only, from a railway bank in 1938 (Tr).
- 92. Geranium pyrenaicum Burm. f. : Mountain Crane's-Bill Quite common in our district: by the railway embankments and along the canal, roadsides, etc. It is replaced northwards by a pale flowered variety.

Page Thirteen

- 93. Geranium molle L. : Dove's-Foot Crane's-Bill Very common throughout the district.
- Geranium pusillum L.: SMALL-FLOWERED CRANE'S-BILL Very occasional; Prime's Farm (Cn).
 - 95. Geranium dissectum L. : CUT-LEAVED CRANE'S-BILL Frequent throughout the district.
 - 96. Geranium robertianum L.: HERB ROBERT One of our commonest plants, occurring throughout the district.
 - 97. Erodium cicutarium agg. : HEMLOCK STORK'S-BILL Occasional only, near Wickham Hall, 1922-1949.

Oxalidaceae

- 98. Oxalis acetosella L.: Common Wood Sorrel Not very frequent but grows sparingly in woods in our district. Hatfield Forest; Birchanger Wood; Bloodhound's Wood; Planting's Wood; Canfield Hart; Alsa Wood in quantity.
- 99. Oxalis corniculata L.: Yellow Wood Sorrel A common weed in gardens from which it sometimes spreads. Not truly wild in our district.

Balsaminaceae

Occasional. In the grounds of the Nursing Home at Stansted, 1940; Stansted old chalk-pit, 1947 (MJ); as a weed in "Meadow Dyke" and other Maze Green Road gardens, 1941 (Cn).

Aquifoliaceae

IOI. Ilex aquifolium L.: HOLLY Very common in all woods in our district.

Celastraceae

102. Euonymus europaeus L. : SPINDLE-TREE Quite frequent in woods and hedges in our district.

Page Fourteen

Rhamnaceae

103. Rhamnus catharticus L. : BUCKTHORN Fairly frequent in hedges and woods in our district. Hatfield Forest.

Aceraceae

Occasional only; not a common tree in our district. Stocking Plantation; by the new water reservoir; Pinchpools, Manuden.

Very common throughout our district both in woods and hedges.

Very frequently planted in the town. Maple Avenue, etc. It is one of the earliest trees to come into flower and leaf in the Spring. The clear gold of the leaves add much to the display of tints in the autumn. A variety, schwedlerii, forms a clump behind the Red White and Blue inn on the Manuden road.

A North American species often planted in gardens in the town.

Little Hadham Road (Kg).

Hippocastanaceae

A very common tree throughout our district both planted in gardens and avenues and woods. The red-flowered Chestnut (A. rubicunda) is also frequent in gardens and by roadsides. A specimen of Aesculus octandra, the Sweet Buck-Eye, grows near the Shell House in Hatfield Forest.

Leguminosae

Common in gardens; also in coppices by the Manuden road near Hazel End, but not truly wild.

110. Genista tinctoria L.: DYER'S GREENWEED
Not frequent, but in scattered places. Brewery Fields for many
years; Tom Wood's Pasture, 1941 (this was ploughed up in 1944);
Golf Course (Hs).

Page Fifteen

- Not a common plant in our district but seen occasionally by roadsides between Stortford and Stansted and also at Takeley, but probably planted in both places. Waste ground near Cherry Gardens.
- 112. Sarothamnus scoparius (L.) Koch: COMMON BROOM
 Not common in our district but in scattered spots. Ugley, outside
 Broome Wood (An); Wallbury Dells (Ad).
- Frequent by roadsides. Prime's Farm; Hadham Road near the Green Man; Farnham gravel-pit; Thorley, by the river; Rickling, etc. Always the spineless variety in our district.
 - II4. Medicago sativa L. : LUCERNE Frequent in cultivation and often as an escape by roadsides.
- 115. Medicago falcata L. : YELLOW SICKLE MEDICK Occasional only. Waterworks Lane, 1942 (Cn); grassfields above Pinchpools, Manuden (An).
 - 116. $Medicago\ lupulina\ L.$: Black Medick (Nonsuch) Plentiful everywhere.
 - 117. Melilotus officinalis (L.) Lam. : Field Melilot Common throughout the district.
- 118. Melilotus alba Medic. : WHITE MELILOT Occasional only. First recorded in 1920 and occasionally since then. Sawbridgeworth, near the railway bridge, 1940 (Ad). Sometimes in cultivated fields with lucerne.
 - 119. Trifolium pratense L. : PURPLE CLOVER Ubiquitous throughout the district.
- 120. Trifolium ochroleucon Huds. : SULPHUR CLOVER Quite frequent by roadsides and hedgerows throughout the district.
- Very occasional, in cultivated grass and clover leys. In a field of rye-grass and clover by Plantings Wood, 1944 (Ad).

Page Sixteen

- 122. Trifolium arvense L.: HARE'S-FOOT TREFOIL
 Very occasional and quite a rarity in our district. Gravel-pit,
 Hallingbury road, by the railway bridge, 1939 (Ad); Stansted road
 (It).
 - 123. Trifolium hybridum L. : ALSIKE CLOVER Very scarce; a rarity in our district. Stansted road (Et).
 - 124. Trifolium repens L. : WHITE (DUTCH) CLOVER Plentiful everywhere.
- 125. Trifolium fragiferum L.: STRAWBERRY-HEADED TREFOIL Occasional in our district. Waterworks Lane, where it has been for many years; Bury Green Lane, 1942 (Ad); near the entrance to Prime's Farm by the small pond, 1946 (Ad).
- 126. Trifolium campestre Schreb.: Hop Trefoil Not uncommon in our district on gravel. Dane's Bridge, 1941 (Ad); Hadham road rifle butts, 1941 (Ad); Bury Green Lane, Isolation Hospital site, 1944 (Ad); Farnham gravel-pit, 1945. And elsewhere
 - 127. Trifolium dubium Sibth. : Lesser Yellow Trefoil Very common throughout our district.
- 128. Lotus corniculatus L.: BIRD'S-FOOT TREFOIL
 One of our commonest plants in fields and by roadsides throughout the district.
- 129. Lotus tenuis Willd.: DECUMBENT BIRD'S-FOOT TREFOIL Found only once, on the site of the old Isolation Hospital, in 1942 and again in the same place in 1946, but not seen since (Ad).
- 130. Lotus uliginosus Schkuhr. : Marsh Bird's-Foot Trefoil Quite frequent in our district. Canfield Hart; Hatfield Forest, etc.
- 131. Robinia pseudo-acacia L.: WHITE ACACIA (FALSE ACACIA)
 A frequent tree in gardens and occasionally found as an escape but not truly wild.
 - 132. Astragalus glycyphyllos L. : SWEET MILK VETCH Very occasional. Quendon road (Cn).
 - 133. Ornithopus perpusillus L. : Common Bird's-Foot Very occasional. Patmore Heath, 1950.

Page Seventeen

- 134. Onobrychis viciifolia Scop. : Sainfoin Commonly cultivated in this district and often found as an escape by roadsides.
- 135. Vicia hirsuta (L.) S. F. Gray : HAIRY VETCH Occasional. Old coach road, Rickling, 1939 (Ad); Farnham chalkpit, 1944.
- 136. Vicia tetrasperma (L.) Schreb. : SLENDER VETCH
 A very common vetch in our district in grassy places and among undergrowth.
 - 137. $Vicia\ cracca\ L.$: Tufted Vetch Ubiquitous in hedgerows.
 - 138. Vicia sepium L. : BUSH VETCH Common throughout the district.
 - 139. $Vicia\ sativa\ L.\ :\ Common\ Vetch\ Common\ everywhere.$
 - 140. Vicia lathyroides L. : Spring Vetch Occasional only. Farnham chalk-pit, 1945.
- 141. Lathyrus aphaca L. : YELLOW VETCHLING Rare. Recorded once only: Maze Green Road, 1922 (Cn), and not seen since.
- 142. Lathyrus nissolia L. : CRIMSON (GRASS) VETCHLING Not uncommon. Golf links (Cn); Spellbrook, 1939; chalk-pit between Little Hadham and Albury, 1940 (Cd).
- 143. Lathyrus pratensis L.: Meadow Vetchling One of the commonest vetchlings in our district, occurring by roadsides everywhere.

Caesalpineae

144. Cercis siliquastrum L. : JUDAS TREE A fine specimen off Parsonage Lane.

Rosaceae

145. Prunus spinosa L. : BLACKTHORN, SLOE Abundant throughout the district.

Page Eighteen

- 146. Prunus insititia L. : BULLACE Abundant throughout the district.
- Fairly common in woods in our district. Hatfield Forest; Bloodhounds Wood; Birchanger Wood; Patmore Wood; Albury, etc.
- Not very common in this district and doubtfully wild. Planted in the Vicarage garden, Bishop's Stortford, and in Sawbridgeworth churchyard.
 - 149. Filipendula ulmaria (L.) Maxim. : Meadowsweet Common in wet places throughout the district.
 - 150. Rubus idaeus L. : Raspberry Hoggett's Wood (Cd).
 - 151. Rubus caesius L. : DEWBERRY Common throughout the distrcit.
 - 152. Rubus fruticosus agg. : Blackberry, Bramble Ubiquitous throughout the district.
 - 153. Geum urbanum L. : Common Avens Common throughout the district.
 - 154. Fragaria vesca L.: WILD STRAWBERRY Very common in woods throughout the district.
- 155. Potentilla sterilis (L.) Garcke. : STRAWBERRY-LEAVED
 CINQUEFOIL
 One of our commonest and earliest Spring flowers.
 - 156. Potentilla verna L. : Spring Cinquefoil.
 Once only—behind the Hospital, 1921 (Cn), and not seen since.
 - 157. Potentilla erecta (L.) Räusch : TORMENTIL Common throughout the district.
 - 158. Potentilla reptans L. : Creeping Cinquefoil Plentiful everywhere.
 - 159. Potentilla anserina L. : SILVERWEED Common in damp places throughout the district.

Page Nineteen

160. Alchemilla arvensis (L.) Scop. : FIELD LADY'S MANTLE (PARSLEY PIERT)

Not very frequent in this district. Sometimes in cornfields;

Not very frequent in this district. Sometimes in cornfields; Broome Wood; a weed in Meadow Dyke garden, Maze Green Road, for many years (Cn).

- 161. Alchemilla vulgaris L., agg. : COMMON LADY'S MANTLE Quite frequent in the district. Hatfield Forest, near the lake; watermeadows near Manuden; near the gamekeeper's cottage, Planting's Wood; Broome Wood, etc.
 - 162. Agrimonia eupatoria L. : COMMON AGRIMONY Common throughout the district; a road- and hedge-side plant.
 - 163. Poterium sanguisorba L. : SALAD BURNET A common plant on the chalk throughout the district.
- 164. Rosa arvensis Huds. : FIELD ROSE
 This white-flowered rose is the commonest in this district in woods and hedges. It is more common than R. canina.
- 165. Rosa canina L.: COMMON DOG ROSE Common in hedgerows throughout the district but not so common as R. arvensis.
- 166. Sorbus aucuparia L.: MOUNTAIN ASH (ROWAN)
 Common in gardens and occasionally as an escape (e.g. Manuden);
 but not truly wild.
 - 167. Sorbus aria agg. : WHITE BEAM Very occasional. Thorley (An).
- Very ocasional. In a hedge at Waterworks Lane (Ad); north side of Canfield Hart wood (Ad).
- 169. Malus pumila Mill.: CRAB APPLE
 Common throughout the district in woods and hedgerows. Common at Ugley (An); Manuden, above Pinchpools (An); Hatfield Forest;
 Wallbury Dells; Stocking Plantation; and other woods.
 - 170. Crataegus oxyacanthoides Thuill. : HAWTHORN A very common tree throughout the district.

Page Twenty

171. Crataegus monogyna Jacq. : CUT-LEAVED HAWTHORN Quite frequent in hedges throughout the district.

Quite a large number of hybrids showing variations of leaf and fruit forms are to be found in the hedges and woods in the district where the Genus is frequent and these need further study. Hybrids of garden origin are found planted as ornamental trees for the sake of their brilliant fruits in the autumn. A fine specimen of C. carrieri Vauv. (C. crus-galli x C. mexicana) is in Maze Green Road.

Saxifragaceae

- 172. Saxifraga granulata L.: Meadow Saxifrage
 Only very occasional in this district. Twyford, 1939 (As); railway
 embankment near the Red White and Blue inn, 1948 (Et).
- 173. Chrysosplenium oppositifolium L.: GOLDEN SAXIFRAGE
 This plant used at one time to grow in Birchanger Wood, but it
 has not been seen there or anywhere else in this district since 1921
 (Cn).

Crassulaceae

- 174. Sedum telephium L.: ORPINE (LIVELONG)
 Rare in this district. Latchmore Bank, in waste ground, 1942;
 cailway bank near Collins Cross, 1943; Broome Wood, 1946.
- 175. Sedum acre L.: BITING STONECROP (WALL PEPPER)
 Doubtfully wild in this district, but often seen in gardens and on the roofs of old cottages.
- 176. Sedum rupestre L. : St. Vincent's Rock Stonecrop Reported once only, from a gravel-pit at Manuden. Probably a garden escape.

Haloragaceae

177. Myriophyllum verticillatum L.: WHORLED WATER MILFOIL Very occasional only. Reported from the canal near Sawbridgeworth (Rs).

Callitrichaceae

178. Callitriche palustris L.: WATER STARWORT Very common in all ponds and streams in the district.

Page Twenty-one

Lythraceae

179. Peplis portula L. : WATER PURSLANE Occasional only.

180. Lythrum salicaria L.: PURPLE LOOSESTRIFE Frequent in wet places throughout the district. In the Town Meads; Hazel End; by the canal, etc.

Onagraceae

181. Chamaenerion angustifolium (L.) Scop. : Rose-Bay
WILLOW-HERB

Very common throughout the district. It appears wherever trees are felled or coppiced. A white-flowered variety appears occasionally.

182. Epilobium hirsutum L. : Great Hairy Willow-Herb Ubiquitous in wet places.

183. Epilobium parviflorum Schreb. : SMALL-FLOWERED Common everywhere.

184. Epilobium montanum L. : Broad-Leaved Willow-Herb Common everywhere.

185. Epilobium roseum Schreb. : PALE WILLOW-HERB Frequent in our district.

These Epilobium in all probability cross, and hybrids showing variations from type can be found.

186. Epilobium tetragonum L. : SQUARE-STALKED WILLOW-HERB Probably quite frequent in this district. Frequent on the banks of the canal (Ad).

187. Epilobium palustre L.: MARSH WILLOW-HERB Occasional. Possibly more frequent but overlooked. Hatfield Forest, in the swamp and watercourse at the head of the lake, 1939 (Et).

188. Circaea lutetiana L. : Enchanter's Nightshade Common throughout the district. Sometimes a troublesome weed in gardens. Very abundant at Hatfield Forest (An).

Cucurbitaceae

189. Bryonia dioica Jacq. : COMMON (WHITE) BRYONY Ubiquitous throughout the district, in hedgerows.

Page Twenty-two

Umbelliferae

190. Hydrocotyle vulgaris L. : Common White-Rot (Marsh Pennywort)
Very occasional; possibly overlooked. Hatfield Forest, 1938.

191. Sanicula europaea L. : WOOD SANICLE Quite common in woods in our district.

192. Conium maculatum L.: HEMLOCK
Not uncommon in the district. Hatfield Forest; Manuden; Rickling road; ditches in Long Meadow; by the River Stort; etc.

193. Bupleurum rotundifolium L.: HARE'S EAR Occasional. Bury Green, 1920 (Cn). Appeared in a garden after a bomb had scattered soil in July 1941 (As).

194. Apium nodiflorum (L.) Lag.: PROCUMBENT MARSHWORT Recorded once only but possibly overlooked in other places. Hat-field Forest, 1929 (Eg).

195. Carum carvi L. : CARAWAY Very occasional. 1927 (Eg).

196. Petroselinum segetum (L.) Koch : CORN PARSLEY Reported once only, 1937 (Eg).

197. Sison amomum L. : BASTARD STONE-PARSLEY (HEDGE SISON)

Frequent in our district. In a ditch, Maze Green Road; Long Meadow; Stocking Plantation, etc.

198. Sium latifolium L.: WATER PARSNIP Occasional. By the Stort, 1941 (Ad). Possibly more common and overlooked.

199. Sium erectum Huds.: NARROW-LEAVED WATER PARSNIP
Frequent in wet places in the district. Hatfield Forest, in the
swamp by the lake; ditches, etc.

200. Aegopodium podagraria L. : GOUT WEED (BISHOP'S WEED)
Ubiquitous throughout the district. Often a troublesome weed in gardens. A common local name for it is Ground Elder.

Page Twenty-three

- 201. Pimpinella saxifraga L.: Common Burnet Saxifrage Rather scarce in our district. In a field by Twyford lock, 1948 (Ad).
- 202. Pimpinella major (L.) Huds. : Burnet Saxifrage Rare. Stocking Plantation; rough ground north of Hallingbury Road near the Sewage Farm, 1942.
 - 203. Conopodium majus (Gouan) Lor. & Barr. : Common Earthnut (Pig Nut)

 Fairly plentiful in the district. Hatfield Forest; in a field below

Fairly plentiful in the district. Hatfield Forest; in a field below the gamekeeper's cottage in Plantings Wood, in quantity (Ad); Wallbury Dells, etc.

- 204. Chaerophyllum temulum L. : CHERVIL
 One of the commonest roadside weeds; ubiquitous throughout
 our district.
 - 205. Scandix pecten-veneris L. : Shepherd's-Needle A very common cornfield weed in this district.
- 206. Anthriscus scandicina (Weber) Mansf. : Beaked Parsley (Burr Chervil)
 Plentiful throughout the district.
 - 207. Oenanthe fistulosa L. : COMMON WATER DROPWORT Occasional. Hatfield Forest, at the west end of the lake, 1946.
 - 208. Aethusa cynapium L. : Fool's Parsley A common weed, plentiful throughout the district.
- 209. Silaum silaus (L.) Schinz & Thell. : Pepper Saxifrage Common throughout the district; a roadside plant. Maple Avenue; plentiful in fields round the College playing-fields.
 - 210. Angelica sylvestris L.: ANGELICA Common in damp places. Hatfield Forest; Birchanger, etc.
- Very plentiful by roadsides throughout the district, especially where the soil is chalky.
- 212. Heracleum sphondylium L. : Cow Parsnip (Hog Weed)
 Ubiquitous throughout the district.

Page Twenty-four

Plentiful throughout the district, especially where the soil is chalky.

Araliaceae

214. Hedera helix L. : Ivy Ubiquitous.

Cornaceae

A very common shrub in this district. (There is a specimen of Cornus mas in a garden at No. 88 Hadham Road.)

Adoxaceae

216. Adoxa moschatellina L.: MOSCHATEL Recorded only once, in 1935, and not seen since.

Caprifoliaceae

- 217. Sambucus nigra L. : COMMON ELDER Plentiful everywhere in our district.
- 218. Viburnum opulus L. : GUELDER ROSE Common in hedgerows throughout the district.
- 219. Viburnum lantana L. : WAYFARING-TREE Common, especially at Manuden and Ugley.
- 220. Symphoricarpus albus (L.) S. F. Blake : SNOWBERRY In woods and shrubberies and often spreads to scrubland in this district. Grows wild in Bloodhounds Wood.
- 221. Lonicera periclymenum L. : Honeysuckle (Woodbine)
 Common throughout the district.

Rubiaceae

222. Galium cruciata (L.) Scop. : CROSSWORT
In one locality only: Hatfield Broad Oak road, between Takeley
station and Canfield Hart wood.

Page Twenty-five

- 223. Galium verum L. : LADY'S BEDSTRAW (YELLOW BEDSTRAW)
 Plentiful everywhere in our district.
 - 224. Galium mollugo L. : HEDGE BEDSTRAW Plentiful throughout the district.
 - 225. Galium palustre L. : MARSH BEDSTRAW Frequent in wet places throughout the district.
 - 226. Galium uliginosum L. : BOG BEDSTRAW Occasional. Hatfield Forest, at the west end of the lake (Ad).
 - 227. Galium aparine L. : Goose-Grass (Cleavers) Ubiquitous.
- 228. Galium tricorne Stokes : ROUGH-FRUITED CORN BEDSTRAW Occasional in hedgerows. Bury Green, 1941 (Cn); bottom of Long Meadow, 1943 (Ad).
 - 229. Asperula odorata L. : WOODRUFF In one locality only: Canfield Hart wood.
 - 230. Sherardia arvensis L.: FIELD MADDER A very common cornfield weed in our district.

Valerianaceae

- 231. Valeriana dioica L. : MARSH VALERIAN

 Not uncommon in wet places in our district. Hatfield Forest; the canal near Spellbrook, etc.
 - 232. Valeriana officinalis L. : Great Wild Valerian Fairly frequent in our district. Hatfield Forest, etc.
- 233. Valerianella locusta L. : CORN SALAD Occasional. Farnham (Cn); railway embankment between the bridges near the Red White and Blue inn (Et).
- 234. Valerianella dentata (L.) Poll. : NARROW-FRUITED CORN
 SALAD
 Fairly common. Farnham (Cn); Much Hadham, 1939; Prime's
 Farm hedgrows, 1941 (Ad).

Page Twenty-six

Dipsaceae

- Very frequent throughout our district. In quantity in Stocking
- 236. Dipsacus pilosus L.: SMALL TEAZEL Occasional only. Much Hadham, 1929; Barwick Ford, 1944 (MJ); in the lane to Farnham gravel-pit. First seen here in 1939, but none in 1940 or 1941. Appeared again in 1942, but absent 1944. Since then II has been seen in 1945, 1947, and 1949. Warren Wood, 1950.
 - 237. Knautia arvensis (L.) Coult. : FIELD SCABIOUS Frequent throughout the district.
- 238. Succisa pratensis Moench: Devil.'s-Bit Scabious
 Not infrequent in restricted localities. Plentiful at one time in
 Tom Wood's Pasture, but this was ploughed in 1944. The plant still
 remains round the margins and in Long Meadow; also round the
 edges of Bloodhounds Wood. Latchmore Bank north of the Hallingbury road, 1944.

Compositae

- 239. Eupatorium cannabinum L. : HEMP AGRIMONY Common in wet places throughout the district.
- 240 Bellis perennis L. : COMMON DAISY Ubiquitous.
- 241. Erigeron canadensis L.: CANADA FLEA-BANE A common weed in waste places. Frequent round about the town.
- 242. Erigeron acris L. : BLUE FLEA-BANE On the verge of Stansted Road, 1950 (Et).
- 243. Filago germanica (L.) L.: COMMON FILAGO Occasional. Near Bloodhounds Wood by the side of a cornfield, 1941 (Fn).
- 244. Filago minima (Sm.) Pers. : LEAST FILAGO Frequent in this district. Maple Avenue, 1938 (Ad); near the river at Manuden, 1939 (Ad); Wallbury Dells, etc.

Page Twenty-seven

- 245. Gnaphalium uliginosum L.: MARSH CUDWEED Reported once only: Latchmore Bank north of the Hallingbury Road, 1944 (Cn).
 - 246. Inula conyza DC. : PLOUGHMAN'S SPIKENARD Stansted old chalk-pit, 1947 (MJ); in Stansted chalk-pit 1951 (Ad).
- 247. Pulicaria dysenterica (L.) Bernh. : COMMON FLEA-BANE Frequent in wet places in our district.
- An introduced weed from Peru. Appeared in a garden in Stansted Road, in 1942 (Hk).
- 249. Bidens cernua L.: Bur Marigold Occasional only. Hatfield Forest by the lake; on the banks of the
- 250. Bidens tripartita L.: TRIFID BUR MARIGOLD
 Not common but seen in several places in our district. Hatfield
 Forest, by the lake; Wallbury Dells, 1937 (Ad); Hazel End, by the
 watercress beds, 1943-5.
 - 251. Achillea millefolium L. : COMMON YARROW Ubiquitous.
 - 252. Anthemis arvensis L. : CORN CHAMOMILE Common throughout the district.
 - 253. Chrysanthemum leucanthemum L. : Ox-EYE DAISY (CORN MARIGOLD)

 A very plentiful plant in cornfields in our district.
 - 254. Matricaria inodora L. : Scentless Mayweed Plentiful everywhere.
- 255. Matricaria matricarioides (Less.) Porter: RAYLESS MAYWEED
 A North American weed, very common by roadsides throughout the district.
 - 256. Tanacetum vulgare L.: TANSY Doubtfully wild in this district. Pledgdon Heath, 1947 (Cn).

Page Twenty-eight

- 257. Artemisia absinthium L.: COMMON WORMWOOD
 One specimen only known in this district which can be considered
 as wild: Elgin Coppice in Hatfield Forest, 1948 (An, Cd).
 - 258. Artemisia vulgaris L. : MUGWORT Very common throughout the district.
- 259. Tussilago farfara L.: COLTSFOOT
 Fairly frequent on railway embankments and by roadsides. Town
 Meads, 1950 (An).
 - 260. Petasites fragrans (Vill.) C. Presl : WINTER HELIOTROPE A garden escape. By the roadside in Maple Avenue.
- 261. Petasites hybridus (L.) Gaertn., Mey & Scherb.: Butterbur Not at all frequent in our district. Fullers End, Elsenham, 1946 (MJ); Cannon Mill Lane, near the railway crossing, 1947 (Et).
 - 262. Senecio vulgaris L. : COMMON GROUNDSEL Ubiquitous.
- 263. Senecio sylvaticus L. : HEATH (MOUNTAIN) GROUNDSEL Comparatively scarce in this district. Brewery fields (Hk); Bloodhounds Wood, 1948 (Et).
- 264. Senecio viscosus L.: STINKING GROUNDSEL
 Reported once only: between the castle and the railway, 1949
 (Kk).
 - 265. Senecio erucifolius L. : HOARY RAGWORT Common throughout the district.
 - 266. Senecio jacobaea L. : COMMON RAGWORT Plentiful everywhere.
 - 267. Senecio aquaticus Hill : Marsh Ragwort Common in wet places by the Stort and canal, etc.
- 268. Carlina vulgaris L.: CARLINE THISTLE
 Very occasional in this district, on chalk. Great Hallingbury, between the church and the lake, 1943 (Ad).
 - 269. Arctium lappa L. : COMMON BURDOCK Ubiquitous.

Page Twenty-nine

- 270. Arctium minus (Hill) Bernh. : Lesser Burdock Quite frequent in our district. Hatfield Forest; Randall's Farm; Takeley, and elsewhere.
 - 271. Carduus crispus L. : WELTED THISTLE Common throughout the district.
 - 272. Carduus nutans L. : MUSK THISTLE Occasional only. Plantings Wood, 1939 (Ad).
- 273. Cirsium vulgare (Savi) Ten. : Spear Plume Thistle Common throughout the district. Very often fasciated, especially in 1937.
 - 274. Cirsium palustre (L.) Scop. : MARSH THISTLE Plentiful everywhere in our district.
 - 275. Cirsium acaulos (L.) Scop. : STEMLESS THISTLE Frequent in this district on chalky soil.
- 276. Cirsium arvense (L.) Scop.: CREEPING (FIELD) THISTLE Ubiquitous. C. arvense var. setosa, a variety without spines on the leaves, appeared in Cricketfield Lane where a pipeline was laid in 1944 and has been there ever since 1949 (Hk).
- 277. Onopordon acanthium L. : Scotch (Cotton) Thistle Rare. Reported once only from a waste dump at the north end of the railway station, 1944 (Pr).
 - 278. Gentaurea nigra L. : COMMON KNAPWEED Plentiful everywhere.
- 279. Centaurea scabiosa L.: Greater Knapweed Common throughout the district, especially on the chalk. Occasionally a white-flowered variety is found. It is frequently parasited by Orobanche major L.
- 280. Centaurea solstitialis L. : Yellow Star Thistle A casual in this country. Reported only once in our district—in a field of lucerne near Farnham chalk-pit, 1949 (Cd).
- 281. Cichorium intybus L.: Succory (Chicory)
 Not uncommon in our district by the edges of fields. Plentiful at
 Bollington Hall (An). Also Long Meadow, Plantings Wood, Bury
 Green Lane, etc.

Page Thirty

- 282. Lapsana communis L. : COMMON NIPPLE-WORT Common throughout the district.
- 283. Picris hieracioides L.: HAWKWEED PICRIS
 Quite frequent in this district. Abundant in the upper playingfields of the College (Cn). Hatfield Forest in quantity, and elsewhere.
 - 284. Picris echioides L. : BRISTLY OX-TONGUE Quite a common plant in our district.
 - 285. Crepis taraxacifolia Thuill. : BEAKED HAWK'S-BEARD Common everywhere by roadsides.
 - 286. Crepis capillaris (L.) Wallr. : SMOOTH HAWK'S-BEARD A very common roadside weed.
 - 287. Hieracium pilosella L. : Mouse-Ear Hawkweed Very common everywhere.
- 288. Hieracium sabauda agg.: SAVOY HAWKWEED In the upper playing-field of the College ten years ago (Cn, 1949); Farnham road, near the gravel-pit, 1944 (Ad).
 - 289. Hypochoeris radicata L. : CAT'S EAR Common everywhere.
 - 290. Leontodon hispidus L. : ROUGH HAWK-BIT Common in grass fields.
 - 291. Leontodon autumnalis L. : AUTUMN HAWK-BIT Common in grass fields.
- 292. Taraxacum officinale Weber: Dandellon Ubiquitous. Has especially fine flowers on the boulder clay soils in this district, of which it is quite a feature.
- 293. Lactuca serriola L.: PRICKLY LETTUCE
 Long Meadow, 1944 (Cn); Thorley Hill, 1944 (Hn); on the north
 side of Canfield Hart wood, 1945 (Ad); in a swamp near the castle
 gardens, 1947 (Ad); seems to be spreading; not uncommon 1949-50.
 - 294. Sonchus oleraceus L. : COMMON SOWTHISTLE Ubiquitous.

Page Thirty-one

- 295. Sonchus asper (L.) Hill: PRICKLY SOWTHISTLE
- 296. Sonchus arvensis L. : CORN SOWTHISTLE Frequent in cornfields.
- 297. Tragopogon pratensis L. : GOAT'S-BEARD Common throughout the district but always the variety with long involucral bracts.
- 298. Tragopogon porrifolius L.: Purple Goat's-Beard (Salsafy) Latchmore Bank, 1948 (MJ). Possibly a garden escape.

Campanulaceae

- 299. Campanula glomerata L.: CLUSTERED BELL-FLOWER
 On the chalk, but not plentiful in this district. Farnham chalkpit; by the footpath in Long Meadow; Bury Green; near Thorley, by the High Wych road, 1944 (Hk).
- 300. Campanula trachelium L.: NETTLE-LEAVED BELL-FLOWER In the lane leading to Farnham chalk-pit; ditch by Stocking Plantation; by the Bourne near Rye Street.
- 301. Campanula rapunculoides L. : CREEPING BELL-FLOWER A garden weed: Meadow Dyke, Maze Green Road. Possibly not truly wild in this district.
- 302. Campanula rotundifolia L. : HAREBELL Hatfield Forest; Rickling Green (Cn); Wallbury Dells; edge of Great Plantings Wood, 1941 (Ad).
 - 303. Specularia hybrida (L.) A. DC. : CORN BELL-FLOWER (VENUS'S LOOKING-GLASS)

 A frequent cornfield weed. Manuden-Rickling old coach road,
- plentiful.

Ericaceae

304. Arbutus unedo L.: STRAWBERRY-TREE A specimen in Bishop's Stortford cemetery.

Page Thirty-two

Monotropaceae

305. Monotropa hypopithys L.: YELLOW BIRD'S-NEST Reported once only—from Stocking Wood, Little Hadham, 1927

Primulaceae

306. Primula vuglaris Huds. : Common Primrose Plentiful in woods in this district.

307. Primula veris L. : CowsLIP Plentiful everywhere in this district.

- 308. Primula elatior (L.) Hill: TRUE OXLIP (BARDFIELD OXLIP)
 Plentiful in woods on the boulder clay in this district. A hybrid
 P. elatior x P. vulgaris occurs in Canfield Hart.
- 309. Lysimachia nummularia L.: Moneywort (Creeping Jenny) Frequent in damp places. Hatfield Forest; Thorley Wood, and elsewhere.
 - 310. Lysimachia nemorum L. : YELLOW PIMPERNEL
- (Wood Loosestrife)

 Birchanger Wood, 1921 (Cn); Canfield Hart, 1946 (Ad); Hatfield

 Forest, 1947 (Ad); Alsa Wood, Ugley, in quantity.
 - 311. Anagallis arvensis L. : SCARLET PIMPERNEL A common cornfield weed.
 - 312. Anagallis foemina Mill. : BLUE PIMPERNEL In a garden at Stansted, 1940.

Oleaceae

313. Fraxinus excelsior L. : Ash A common tree throughout the district.

314. Ligustrum vulgare L.: PRIVET
Frequent in woods, where it has probably spread from gardens by bird-carried seed.

Apocynaceae

315. Vinca minor L.: LESSER PERIWINKLE
At the Much Hadham road end of Bury Green Lane and in the opposite wood.

Page Thirty-three

Gentianaceae

- 316. Blackstonia perfoliata (L.) Huds. : Perfoliate Yellow-Wort
- On chalk. Rifle butts by Wickham Hall (now demolished); north side of Canfield Hart wood.
- 317. Centaurium umbellatum Gilib. : Centaury
 Frequent in our district, especially on chalky soils. A whiteflowered variety is not uncommon.

Boraginaceae

- 318. Cynoglossum officinale L. : Common Hound's-Tongue Plentiful at Hatfield Forest; also at Canfield Hart.
- 319. Symphytum officinale L.: COMMON COMFREY Plentiful by the river and canal and in wet places. Hatfield Forest, outside Beggar's Hall.
- 320. Anchusa sempervirens L.: EVERGREEN ALKANET Bishop's Stortford, 1926 (Cn), 1938 (Ad); Bury Lodge; Takeley, rear the A.A. box, 1941 (Ad).
 - 321. Myosotis scorpioides L. : Forget-Me-Not Frequent in all wet places.
 - 322. Myosotis arvensis (L.) Hill: FIELD SCORPION-GRASS Common everywhere throughout the district.
- 323. Myosotis collina Hoffm.: EARLY SCORPION-GRASS
 This small Myosotis is frequent at Hatfield Forest on the mounds
 made by the ant Ananthomyops flavus. Also found on the railway
 embankment between the bridges north of the Red White and Blue
 inn, 1948 (Et).
- 324. Myosotis versicolor Sm.: Yellow and Blue Forget-Me-Not We have only one record: Thorley, 1949 (Et).
 - 325. Lithospermum officinale L. : COMMON GROMWELL Recorded only at Hatfield Forest.
 - 326. Lithospermum arvense L. : CORN GROMWELL (BASTARD ALKANET)
 A very common cornfield weed throughout the district.

Page Thirty-four

Convolvulaceae

- 327. Calystegia sepium (L.) R. Br. : Greater Bindweed Plentiful everywhere and often a troublesome weed in gardens.
- 328. Convolvulus arvensis L. : LESSER BINDWEED Common throughout the district; a roadside weed.

Solanaceae

- 329. Solanum dulcamara L. : WOODY NIGHTSHADE (BITTERSWEET)
 Common throughout the district.
 - 330. Solanum nigrum L. : BLACK NIGHTSHADE A common weed everywhere.
- 331. Solanum sisymbrifolium Lam. An introduced plant which appeared in a field of flax on Pocock's Farm in 1941. The flax seed probably came from Brazil.
 - 332. Lycium chinense Mill. : TEA-PLANT (TEA-TREE) Frequently cultivated as a hedge plant in this district.
- 333. Atropa bella-donna L. : DEADLY NIGHTSHADE (DWALE) Stansted chalk-pit; Manuden.
 - 334. Datura stramonium L. : COMMON THORN-APPLE Quite frequent in waste places and near rubbish heaps.
- 335. Hyoscyamus niger L.: Henbane
 Fairly common in our district. The College vegetable garden;
 Convent playing-fields; Hockerill College, in a filled-in bomb crater,
 1942; Farnham gravel-pit; and elsewhere.

Scrophulariaceae

- 336. Verbascum thapsus L.: Great Mullein Hatfield Forest, in Elgin Coppice, 1948-50 (An), and Spittlemore Coppice, 1947; Farnham chalk-pit.
- 337. Verbascum nigrum L. : DARK MULLEIN (BLACK MULLEIN) Occasional only.
 - 338. Verbascum blattaria L. : MOTH MULLEIN Occasional; usually a garden escape.

Page Thirty-five

- 339. Linaria cymbalaria (L.) Mill. : IVY-LEAVED TOAD-FLAX Frequent on walls in the town. Bury Lodge, Takeley, 1941 (Ad).
- 340. Linaria elatine (L.) Mill. : SHARP-POINTED FLUELLIN Occasional only. In a cornfield on the east side of Hatfield Forest, not far from Shell House, 1946 (Pr).
 - 341. Linaria spuria (L.) Mill. : MALE FLUELLIN (ROUND-LEAVED TOAD-FLAX)
 A fairly frequent cornfield weed in our district.
 - 342. Linaria purpurea (L.) Mill. : PURPLE TOAD-FLAX)
 On waste land by the river. Probably a garden escape.
 - 343. Linaria vulgaris Mill. : YELLOW TOAD-FLAX Not uncommon by roadsides and on railway embankments.
- 344. Linaria minor (L.) Desf.: Lesser Toad-Flax Long Meadow, 1922 (Cn); Stane Street halt, 1946-50 (Ad); Thorley Wash, in ballast on the railway line, 1948-49.
- 345. Scrophularia aquatica L.: WATER FIGWORT
 Frequent in wet places, ditches, and along the river-bank above
 the town.
 - 346. Scrophularia nodosa L. : KNOTTED FIGWORT Common throughout the district.
 - 347. Scrophularia vernalis L.: Yellow Figwort Recorded once only, at Debden in 1939 (Cn).
 - 348. Digitalis purpurea L. : Foxglove Ugley and Quendon woods (An), but rare in our district.
 - 349. Veronica hederifolia L. : IVY-LEAVED SPEEDWELL A very common garden weed.
 - 350. Veronica persica Poir. : BUXBAUM'S SPEEDWELL Plentiful everywhere.
- 351. Veronica arvensis L.: WALL SPEEDWELL Fairly common on walls and in waste places and woods in the district.

Page Thirty-six

- 352. Veronica serpyllifolia L. : THYME-LEAVED SPEEDWELL Common everywhere.
- 353. Veronica officinalis L.: COMMON SPEEDWELL
 Not so frequent as some other Veronicas in this district but
 occurs in woods—Hatfield Forest, Bloodhound's Wood, Canfield Hart
 and others.
 - 354. Veronica chamaedrys L. : GERMANDER SPEEDWELL Plentiful everywhere.
 - 355. Veronica scutellata L. : MARSH SPEEDWELL Woodside Green (Cn); Patmore Heath, 1950.
 - 356. Veronica montana L.: MOUNTAIN SPEEDWELL Alsa Wood, Ugley, 1950. Also in other woods, 1950.
- 357. Veronica anagallis-acquatica L.: WATER SPEEDWELL In ditches and wet places. Waterworks Lane, 1941; Thorley, near the church, 1944; Hatfield Forest, on the lake margins, 1946.
 - 358. Veronica beccabunga L. : BROOKLIME Frequent in wet places.
- 359. Euphrasia officinalis agg.: EYEBRIGHT
 Not uncommon, especially on chalky soil. Hatfield Forest; Woodside Green; Farnham chalk-pit; Takeley; Randall's Farm, and elsewhere.
 - 360. Odontites rubra Gilib. : RED BARTSIA Common throughout the district.
 - 361. Rhinanthus minor agg. : COMMON YELLOW RATTLE A hayfield weed. Spellbrook watermeadows.
 - 362. Melampyrum cristatum L. : CRESTED COW-WHEAT Manuden, 1949 (Hk and Et).
 - 363. Melampyrum arvense L.: PURPLE COW-WHEAT Long Meadow, in a barley field, 1920 (Cn). Not recorded since.
- 364. Melampyrum pratense L. : Common Yellow Cow-Wheat Occasional. Wallbury Dells; Weston Green; Little Hadham (Cn).

Page Thirty-seven

Orobanchaceae

- 365. Orobanche elatior Sutton: Tall Broom-Rape Frequent in this district, parasitic on Centaurea scabiosa L. In quantity in Farnham gravel-pit.
- 366. Orobanche apiculata Wallr.: SMALL BROOM-RAPE Stortford Park and Hyde Hall cricket-ground (Cn); Manuden, 1942 (Hk); near the lake, Great Hallingbury, 1943; Birchanger, in abundance, 1949 (Et). Parasitic on clover.

Verbenaceae

367. Verbena officinalis L. : Vervain Bedlar's Green, 1943 (Hs).

Labiatae

- 368. Mentha longifolia (L.) Huds.: Horse Mint Gaston Green 1948 (Rs); between the Castle and the railway, 1949 (Kk).
 - 369. Mentha spicata L. : Spearmint Between the Castle and the railway, 1949 (Kk).
 - 370. Mentha aquatica L. : WATER MINT Very common in ditches and wet places.
- 371. Mentha gracilis Sole: Whorled Mint This plant is probably of hybrid origin. Bloodhounds Wood, 1940 (Cn); Stocking Plantation, 1945; and elsewhere.
- 372. Mentha arvensis L.: CORN MINT Frequent in cornfields. Manuden (Ad); Randall's Farm, Takeley, 1940 (Ad); and elsewhere.
- 373. Lycopus europaeus L. : GIPSY-WORT
 Common by the side of the canal and elsewhere in wet places.
 Hatfield Forest.
- 374. Origanum vulgare L.: Marjoram Farnham gravel-pit, in quantity; Rickling Green, old coach road; Much Hadham road, 1944 (Ad).
 - 375. Thymus serpyllum agg. : WILD THYME Farnham chalk-pit (An); Hatfield Forest; Patmore Heath.

Page Thirty-eight

- 376. Acinos arvensis (Lam.) Dandy: Basil Thyme Fairly common in hedgerows.
- 377. Calamintha ascendens Jord.: Common Calamint Farnham gravel-pit; Hatfield Forest; and elsewhere. Common.
- 378. Salvia verbenaca L.sec.Pugsl. : CLARY (WILD SAGE) Scarce. Puckeridge, 1949; Stansted road, 1946.
- 379. Nepeta cataria L.: CATMINT Farnham Lane, 1948 (Et), and again in 1950.
- 380. Glechoma hederacea L. : GROUND IVY A common weed everywhere.
- 331. Scutellaria galericulata L. : Common Skullcap On the river and canal banks.
- 382. Prunella vulgaris L. : COMMON SELF-HEAL Plentiful everywhere.
- 383. Stachys officinalis (L.) Trev. : Wood Betony Tom Wood's Pasture (since ploughed); Long Meadow; Wallbury Dells; Birchanger; Hockerill field-path.
 - 384. Stachys palustris L. : MARSH WOUNDWORT By the river and canal.
 - 385. Stachys sylvatica L. : HEDGE WOUNDWORT Plentiful everywhere.
- 386. Stachys annua (L.) L. : PALE ANNUAL WOUNDWORT Appeared as a weed in Mr. Sutton's garden, Maze Green Road, in 1940.
 - 387. Galeopsis ladanum L.: HEMP-NETTLE In the Rickling-Berden old coach road.
 - 388. Galeopsis angustifolia Hoffm.: RED HEMP-NETTLE Little Hadham Farm, 1941, and elsewhere. Not uncommon.
- 389. Galeopsis tetrahit L.: Common Hemp-Nettle Frequent. Wallbury Dells, 1939 (Ad); hedgerows at Prime's Farm, and elsewhere.

Page Thirty-nine

- 390. Lamium amplexicaule L. : HENBIT Farnham gravel-pit, 1942 (MJ).
- Lamium hybridum Vill. : Hybrid Red Dead Nettle Maze Green Road, 1949 (Ad and Kk).
- 392. Lamium purpureum L. : RED DEAD NETTLE Ubiquitous.
- 393. Lamium album L. : WHITE DEAD NETTLE Ubiquitous.
- 394. Galeobdolon luteum Huds.: Yellow Archangel Frequent in this district. Long Meadow; Bury Green; Hadham road; Takeley road, and elsewhere.
 - 395. Ballota nigra L. : BLACK HOREHOUND A common roadside and hedgerow plant.
- 396. Ajuga reptans L.: Common Bugle
 Very common everywhere. A variety with pink flowers occurs in
 Bloodhounds Wood and one with white flowers in Elgin Coppice,
 Hatfield Forest, and in Canfield Hart.

Plantaginaceae

- 397. Plantago lanceolata L. : RIBWORT PLANTAIN Ubiquitous.
- 398. Plantago media L. : HOARY PLANTAIN Ubiquitous.
- 399. Plantago major L. : Greater Plantain Ubiquitous.

Illecebraceae

400. Scleranthus annuus L. : Annual Knawel Cornfields near Hallingbury, 1942.

Amaranthaceae

401. Amaranthus retroflexus L.
An introduced plant which appeared in a field of lucerne on Wickham Hall farm in 1949 (Cd).

Page Forty

Chenopodiaceae

- 402. Chenopodium polyspermum L. : MANY-SEEDED GOOSEFOOT A very common weed in our district.
 - 403. Chenopodium album L. : White Goosefoot Common everywhere.
 - 404. Chenopodium urbicum L.: UPRIGHT GOOSEFOOT Hatfield Forest, by the lake, 1938; Manuden.
 - 405. Chenopodium rubrum L. : RED GOOSEFOOT Common everywhere.
- 406. Chenopodium bonus-henricus L. : GOOD KING HENRY Long Meadow; the canal, near the gasworks, 1940 (Ad); Manuden (An); and elsewhere.
- 407. Atriplex patula L.: COMMON ORACH Manuden-Rickling road, 1938 (Ad); Thorley, by the river, 1949 (Et); Woodside Green, 1949 (Et).

Polygonaceae

- 408. Polygonum convolvulus L. : BLACK BINDWEED A common cornfield weed in our district.
- 409. Polygonum dumetorum L. : COPSE BUCKWHEAT A weed in the garden of Meadow Dyke, Maze Green Road (Cn).
- 410. Polygonum aviculare agg. : COMMON KNOTGRASS
 Common on waste ground in the Windhill district, by the gasworks, etc.
 - 411. Polygonum hydropiper L. : WATER PEPPER Hatfield Forest, by the lake; waste ground near the canal.
 - 412. Polygonum persicaria L. : Spotted Persicaria Hatfield Forest, by the lake; waste ground by the gasworks (Ad).
- 413. Polygonum lapathifolium L. : PALE-FLOWERED PERSICARIA Waste ground by the gasworks (Ad); Cricket Field, 1950 (Hk).

Page Forty-one

- 414. Polygonum amphibium L. : Amphibious Persicaria Waste ground by the gasworks (Ad); Twyford lock (Cn).
- 415. Polygonum bistorta L.: Common Bistort (Snakewort)
 Maze Green Road, in a field since built upon, 1937; Sewage Farm.
- 416. Rumex conglomeratus Murr. : Sharp Dock Bollington Hall farm, near Manuden, 1949 (Et); Stocking Plantation, 1949 (Et).
 - 417. Rumex obtusifolius L. : BROAD-LEAVED DOCK Common everywhere.
 - 418. Rumex crispus L. : CURLED DOCK Harvey's Farm, 1939 (Ad).
 - 419. Rumex sanguineus L. : RED-VEINED DOCK Woodside Green, 1949 (Et).
 - 420. Rumex hydrolapathum Huds. : Great Water Dock On the canal banks.
- 421. Rumex acetosella L. : Sheep's Sorrel Common. Patmore Heath (An); Wallbury Dells; Hatfield Forest; and elsewhere.
 - 422. Rumex acetosa L. : Common Sorrel Common everywhere.

Thymelaeaceae

423. Daphne laureola L.: Spurge Laurel Bury Green; Coppice at back of Thorley road (As); in quantity in the wood between Stocking Plantation and Great Hadham road; Hoggett's Wood.

Loranthaceae

424. Viscum album L.: MISTLETOE
Plentiful at Hatfield Forest on hawthorn and other trees; several
colonies on Aesculus octandra near the Shell House.

Page Forty-two

Euphorbiaceae

- 425. Euphorbia peplis L. : PURPLE SPURGE Common everywhere.
- 426. Euphorbia platyphyllos L.: BROAD-LEAVED WARTED SPURGE Bury Green Lane, 1944 (Cn); edge of Canfield Hart, 1945 (Ad).
 - 427. Euphorbia esula L. : LEAFY SPURGE Wallbury Dells; Stane Street Halt, by the railway.
 - 428. Euphorbia exigua L. : DWARF SPURGE A common comfield weed in our district.
 - 429. Euphorbia lathyrus L. : CAPER SPURGE Common in gardens as a weed; probably not truly wild.
 - 430. Mercurialis perennis L. : Dog's MERCURY Ubiquitous.
- 431. Mercurialis annua L.: Annual Mercury
 Thornfield Road, 1945 (MJ); as a garden weed at 42 Hadham
 Road, 1949 (Cn).

Buxaceae

432. Buxus sempervirens L. : Box Hatfield Forest; doubtfully wild.

Ulmaceae

- 433. Ulmus glabra Huds. : WYCH ELM Frequent in this district.
- 434. Ulmus procera Salisb. : Common Elm Frequent in this district, but many are dying from elm disease.

Cannabinaceae

435. Humulus lupulus L. : Hop
Common in hedges. Probably escape from cultivation; but no
hops are now grown in this district.

Page Forty-three

Urticaceae

436. Urtica dioica L. : STINGING NETTLE Ubiquitous.

437. Urtica urens L. : SMALL NETTLE Stansted chalk-pit, 1937 (Ad); Bury Green Lane, 1946 (Ad) Alsa Wood, near Ugley Green, 1950; Gilston Park, 1950.

438. Parietaria diffusa Mert. & Koch. : Pellitory-of-the-Wali On the bridge over the stream between Spellbrook and the railway crossing; on the wall of a house in Manuden.

Juglandaceae

439. Juglans regia L. : WALNUT A fairly common tree in this district.

440. Juglans nigra L. : BLACK WALNUT Hatfield Forest, in the chestnut grove at Warren House.

Betulaceae

441. Betula pendula Roth: Common Birch (Silver Birch)
A common tree in this district.

442. Alnus glutinosa (L.) Gaertn. : Common Alder Hatfield Forest; Stocking Plantation; by the Dunmow road near Thremhall Priory. Occasional in the N.W. area.

Corylaceae

443. Carpinus betulus L. : HORNBEAM With the oak the commonest tree in our district.

444. Corylus avellana L.: HAZEL
The common undergrowth in the woods in this district and in hedgerows.

Fagaceae

The most prominent tree in this district, all of the pedunculate kind.

Page Forty-four

446. Quercus cerris L.: TURKEY OAK
A large specimen at Great Hallingbury Church; Hatfield Forest,
in Spittlemore Coppice.

447. Quercus ilex L. : HOLM OAK Hazel End; Windhill monastery garden.

448. Quercus suber L. : CORK OAK One historic specimen at Wallbury Dells.

449. Castanea sativa Mill. : SPANISH CHESTNUT Frequent in woods in this district. Hatfield Forest, etc.

450. Fagus sylvatica L. : BEECH Occasional only in this district. Hatfield Forest; Ugley; and elsewhere.

Salicaceae

451. Salix fragilis L.: CRACK WILLOW

By the pond near the entrance to Prime's farm, and elsewhere.

452. Salix alba L.: COMMON WHITE WILLOW Frequent in this district. A variety, S. vitellina, has been planted by the roadside at Takeley.

453. Salix viminalis L.: OSIER
Hazel End; by the River Stort and canal; near the Castle gardens.
Often cultivated.

454. Salix caprea L. : GOAT WILLOW Frequent throughout the district, but rather rare north of Pinchpools (Manuden).

455. Salix aurita L. : ROUND-LEAVED WILLOW Frequent.

456. Salix atrocinerea Brot. : GREY SALLOW Frequent.

457. Salix babylonica L.: Weeping Willow In gardens at Hassobury; Great Hallingbury, near the church.

Page Forty-five

458. Populus alba L. : White Poplar (Abele) Frequent in this district.

459. Populus tremula L. : ASPEN
Plentiful, sometimes abundant, in most woods in the district.
Hatfield Forest; Bloodhounds Wood; Hoggett's Wood, etc.

460. Populus nigra L. : BLACK POPLAR Frequent; often planted.

461. Populus fastigiata L.: LOMBARDY POPLAR Manuden, in the glebe (planted); occasional elsewhere.

462. Populus balsamifera L. : Balsam (Canadian) Poplar Garden of White Lodge; College grounds.

463. Populus trichocarpa L. A small plantation near the Reindeer Inn at Spellbrook. A native of western North America.

Ceratophyllaceae

464. Ceratophyllum demersum L.: Common Hornwort In the canal (Ad); in the lake at Hatfield Forest (Ad).

MONOCOTYLEDONS

Hydrocharitaceae

465. Elodea canadensis Michx.: WATER THYME In the canal; in a stream by Stocking Plantation; in the lake at Hatfield Forest.

Orchidaceae

466. Neottia nidus-avis (L.) L. C. Rich. : BIRD'S-NEST ORCHIS "I have notes of this in various years up to 1949 from Long Meadow and Stansted" (Cn).

467. Listera ovata (L.) R. Br. : TWAYBLADE
Not uncommon in woods in this district: Hatfield Forest;
Canfield Hart; Planting's Wood; Thorley Wood; etc.

Page Forty-six

- 468. Anacamptis pyramidalis (L.) L. C. Rich. : PYRAMIDAL ORCHIS
- College Upper Field (Cn); between Great Hallingbury church and the lake, 1943; Hatfield Forest, 1946; gravel-pit at Manuden, 1949 (Et).
- 469. Orchis morio L.: Green-Winged Orchis "Used to be abundant in and just beyond Long Meadow but has not been seen there for some years" (Cn).
 - 470. Orchis mascula (L.) L. : EARLY PURPLE ORCHIS Frequent in all woods in our district.
- 471. Orchis latifolia L. sec. Pugsl.: MARSH ORCHIS Frequent in wet places in this district. A complete series of colours can be found. This species now includes what was sometimes called Spotted Orchis, the O. maculata of B. & H.
- 472. Ophrys apifera Huds. : BEE ORCHIS
 Not uncommon in this district but very variable, being plentiful
 in some years (1937 particularly) and scarce in others. Golf course;
 College Upper Field; Farnham gravel-pit; Wickham Hall rifle-butts;
 Canfield Hart, N.E. side; Hatfield Forest.
- 473. Ophrys insectifera L.: FLY ORCHIS
 "One specimen found in Thorley Wood in 1925" (Cn). Not seen since.
- 474. Gymnadenia conopsea (L.) R. Br. : Fragrant Orchis Spellbrook, 1939.
 - 475. Coeloglossum viride (L.) Hartm. : Green Habenaria (Frog Orchis)
- "Used to be frequent in Long Meadow but not seen there for many years" (Cn).
- 476. Platanthera bifolia (L.) L. C. Rich.: Butterfly Orchis Sparingly in woods in this district. Hatfield Forest; Canfield Hart; Planting's Wood; Thorley Wood; Battles Wood, Manuden, 1942 (Hk).

Iridaceae

477. Iris foetidissima L. : STINKING IRIS (GLADDON) Canfield Hart; Birchanger Wood.

Page Forty-seven

478. Iris pseudacorus L. : Yellow Flag Iris Frequent in wet places by the river and canal.

Amaryllidaceae

479. Narcissus pseudo-narcissus L.: WILD DAFFODIL Farnham; Bloodhounds Wood; Crouch Hill Wood, Manuden. Formerly (1938) common in Beggar's Hall Coppice, Hatfield Forest, but since picked out of existence (An).

480. Allium vineale L. : CROW GARLIC Frequent as a weed in cornfields and by roadsides.

481. Allium ursinum L.: RAMSONS
Rare in this district. "In a swampy field by the roadside at Little
Hallingbury, 1948" (Rs); Battles Wood, Manuden, 1950; Patmore
Heath Wood, 1950.

Dioscoreaceae

482. $Tamus\ communis\ L.$: Black Bryony Frequent in hedgerows.

Liliaceae

483. Scilla non-scripta (L.) Hoffm. & Link : WILD HYACINTH (BLUEBELL)
Common in all our woods.

484. Ornithogalum umbellatum L.: STAR OF BETHLEHEM Occasional; doubtfully wild. By the side of the Manuden-Rickling road, near Rickling, 1936 (An).

485. Paris quadrifolia L. : Herb Paris To be found in most woods in our district.

Juncaceae

486. Juncus inflexus L. : HARD RUSH Hatfield Forest. Frequent in wet places.

487. Juncus effusus L. : Soft Rush Frequent by the river and canal.

Page Forty-eight

- 488. Juncus articulatus L. : COMMON JOINTED RUSH In the town (As).
- 489. Luzula pilosa (L.) Willd. : HAIRY WOOD-RUSH Birchanger wood; Bloodhounds Wood; Hatfield Forest.
- 490. Luzula campestris (L.) DC. : FIELD WOOD-RUSH Common everywhere in dry open fields.

Typhaceae

491. Typha latifolia L.: GREAT REED-MACE
By the river and canal; Sewage Farm, 1949; Hatfield Forest, by
the lake.

Sparganiaceae

- 492. Sparganium erectum L.: Branched Bur-Reed Frequent in wet places in our district. Hatfield Forest.
- 493. Sparganium simplex Huds. : SIMPLE BUR-REED By the canal, 1940 (Ad); in the moat between the Castle gardens and the Causeway; near Sawbridgeworth station, 1948 (Rs).

Araceae

494. Arum maculatum L. : Cuckoo-Pint (Lords And Ladies)
Ubiquitous.

Lemnaceae

495. Lemna trisulca L.: IVY-LEAVED DUCKWEED Hatfield Forest, in the swamp at the N.W. end of the lake.

496. Lemna minor L. : Lesser Duckweed Common everywhere.

Alismataceae

- 497. Alisma plantago-aquatica L. : Great Water Plantain Plentiful by the river and canal. Hatfield Forest, by the lake.
 - 498. Sagittaria sagittifolia L.: COMMON ARROWHEAD Common in the river and canal and in streams in this district.

Page Forty-nine

Butomaceae

499. Butomus umbellatus L.: FLOWERING RUSH By the canal near Sawbridgeworth, 1938; in a pond at Wickham Hall farm, 1949 (Cd).

Potamogetonaceae

- 500. Potamogeton natans L. : BROAD-LEAVED POND-WEED Common in ponds and in the canal.
- 501. Potamogeton perfoliatus L. : Perfoliate Pond-Weed In the canal, 1938.
 - 502. Potamogeton crispus L. : CURLY POND-WEED In the canal, 1938.
 - 503. Potamogeton pectinatus L. : Fennel Pond-Weed In the canal, 1938.
- 504. Potamogeton densus L. : Opposite-Leaved Pond-Weed In the lake at Hatfield Forest; in water-cress beds near Pinchpools.

Cyperaceae

- 505. Scirpus lacustris L. : Common Bull-Rush (Bulrush) By the river and canal.
 - 506. Scirpus sylvaticus L. : WOOD CLUB-RUSH Once only : at Sheering in 1947.
- 507. Blysmus compressus (L.) Link: BROAD-LEAVED BLYSMUS Hatfield Forest, at the south end of the lake.
- 508. Cladium mariscus (L.) Pohl: Fen Sedge Hatfield Forest, at the N.W. end of the lake, 1944. "By 1949 it had spread into the fen area, and as this silted up it increased there to a great extent". (An).
 - 509. Carex pulicaris L. : FLEA SEDGE Common everywhere.
 - 510. Carex disticha Huds. : Soft Brown Sedge Hatfield Forest, 1945 (As).

Page Fifty

- 511. Carex paniculata L. : PANICLED SEDGE Hatfield Forest.
- 512. Carex vulpina L.: Fox Sedge
 Common in wet places by the canal; Hatfield Forest by the lake, and elsewhere.
 - 513. Carex spicata Huds. : PRICKLY SEDGE By the river and canal; Hatfield Forest (As).
 - 514. Carex echinata Murr. : LITTLE PRICKLY SEDGE Common everywhere.
 - 515. Carex remota L. : REMOTE SEDGE Common everywhere.
- 516. Carex nigra (L.) Reichard : BLACK SEDGE
 Hatfield Forest; Thorley, meadows by the railway, 1940 (As);
 Bailey Hills woods, 1944; Patmore Heath, 1950 (Ad).
 - 517. Carex flacca Schreb. : GLAUCOUS SEDGE Common everywhere.
- 518. Carex pendula L.: PENDULOUS SEDGE Hatfield Forest; Bury Green Lane, 1939 (As); frequent in this district.
 - 519. Carex sylvatica Huds. : Wood Sedge Frequent in woods in our district.
 - 520. Carex hirta L. : HAIRY SEDGE Hatfield Forest; Patmore Heath, 1950 (Ad).
 - 521. Carex ovalis Good. : OVAL SEDGE Patmore Heath, 1950 (Ad).
- 522. Carex pseudocyperus L.: CYPERUS-LIKE SEDGE Hatfield Forest, at the north end of the lake (As); Plantings Wood (As).
 - 523. Carex acutiformis Ehrh.: MARSH SEDGE By the River Stort, 1940; town meads, 1947 (As); Hatfield Forest.

Page Fifty-one

524. Carex vesicaria L. : BLADDER SEDGE Hatfield Forest, by the side of the lake, 1947.

Gramineae

- 525. Setaria viridis (L.) Beauv. : Green Bristle Grass An introduced species. Near the cricket ground, 1949 (Kk); in a field of lucerne, Wickham Hall farm, 1949 (Cd).
 - 526. Phalaris canariensis L. : Canary Grass Occasional. Generally in gardens.
- 527. Phalaris arundinacea L.: COMMON REED-GRASS Abundant in the streams round the car park; town meads; by the gasworks, etc.
 - 528. Anthoxanthum odoratum L. : Vernal Grass Hockerill meadows, and elsewhere.
- 529. Alopecurus myosuroides Huds. : SLENDER FOX-TAIL GRASS On the Manuden-Rickling road, 1938.
 - 530. Alopecurus pratensis L. : MEADOW FOX-TAIL Common everywhere.
 - 531. Milium effusum L. : Spreading Millet Occasional. In the town, 1940.
- 532. Phleum pratense L. : COMMON CAT'S-TAIL (TIMOTHY)
 GRASS
- 533. Agrostis canina L. : Brown Bent-Grass
 Birchanger Wood, 1939; Broome Wood, Ugley, 1946; Hatfield
 Forest, 1946.
 - 534. Agrostis stolonifera L. : Marsh Bent-Grass Abundant in Hatfield Forest.
- 535. Calamagrostis epigejos (L.) Roth : Wood Small-Reed (Bush-Grass)
 In open spaces in thickets. Hatfield Forest; Alsa Wood, Stansted, 1950 (Ad).

Page Fifty-two

- 536. Aira caryophyllea L. : SILVERY HAIR-GRASS Patmore Heath, 1950 (Ad).
- 537. Aira praecox L. : EARLY HAIR-GRASS Patmore Heath, 1950 (Ad).
- 538. Deschampsia caespitosa (L.) Beauv. : Tufted Hair-Grass Patmore Heath, 1950 (Ad).
- 539. Deschampsia flexuosa (L.) Trin. : WAVED HAIR-GRASS On hedge banks. Seldom flowers.
 - 540. Holcus mollis L. : Creeping Soft-Grass In the town, 1939 (As).
 - 541. Holcus lanatus L. : Meadow Soft-Grass (Yorkshire Fog)
 Common everywhere.
 - 542. Avena fatua L.: WILD OAT
 By the canal towing-path near Sawbridgeworth, 1948 (Rs).
 - 543. Arrhenatherum elatius (L.) J. & C. Presl. : FALSE OAT Common everywhere.
- 544. Phragmites communis Trin.: Common REED Common in wet places by the river and canal. Hatfield Forest, by the lake.
 - 545. Cynosurus cristatus L. : CRESTED DOG'S-TAIL GRASS Hockerill meadows, 1939; Patmore Heath, 1950 (Ad).
 - 546. Koeleria gracilis Pers. : Crested Keoleria Patmore Heath, 1950 (Ad).
- By the River Stort, 1940 (As). Found in much the same place as recorded in the Flora Hertfordensis. Also on dredged mud by the river at South Mill, 1940 (As); between the Castle and the railway at Stortford, 1949 (Pr and Kk).
 - 548. Melica uniflora Retz. : Wood MELICK Occasional.

Page Fifty-three

- 549. Dactylis glomerata L. : Cock's-Foot Grass Common everywhere.
- 550. Briza media L. : COMMON QUAKING GRASS Common. Hatfield Forest, etc.
- 551. Poa annua L. : Annual Meadow-Grass Universal.
- 552. Poa compressa L. : FLAT-STEMMED MEADOW-GRASS Between the Castle and the railway, Bishop's Stortford, 1949 (Kk).
 - 553. Poa pratensis agg. : SMOOTH MEADOW-GRASS Common everywhere.
 - 554. Poa trivialis L. : ROUGHISH MEADOW-GRASS Occasional.
- 555. Glyceria fluitans (L.) R. Br. : FLOTE-GRASS Occasional.
- 556. *Glyceria maxima* (Hartm.) Holmb. : Reed Meadow-Grass By the river and canal.
- 557. Festuca pratensis Huds. : Meadow Fescue-Grass College meadows, 1939; Hockerill meadows; on hedge banks in Maple Avenue; the town meads.
 - 558. Bromus ramosus Huds. : HAIRY WOOD BROME-GRASS In the town, 1939 (As); Sawbridgeworth.
 - 559. Bromus sterilis L. : Barren Brome-Grass Common everywhere.
- 560. Brachypodium sylvaticum (Huds.) Beauv. : SLENDER FALSE
 BROME-GRASS
 Occasional.
 - 561. Lolium perenne L. : PERENNIAL RYE-GRASS Universal.
 - 562. Lolium temulentum L. : DARNEL Common everywhere.

Page Fifty-four

- 563. Agropyron repens (L.) Beauv. : COUCH-GRASS (QUITCH, TWITCH)
 Universal.
- 564. Nardus stricta L. : MAT-GRASS Patmore Heath, 1950 (Ad).
- 565. Hordeum nodosum L. : Meadow Barley Occasional.
- 566. Hordeum murinum L. : WALL BARLEY Common everywhere.

CONIFERAE

A number of species in this Natural Order occur in our district. Most of these have been introduced and have been planted for ornamental purposes in gardens in and around the town and in parks like Hatfield Forest.

The following list has been compiled by Mr. H. J. Holmes of our Society.

Cupressaceae

567. Juniperus communis L. : COMMON JUNIPER

Taxaceae

568. Taxus baccata L.: Common Yew
Frequent in churchyards throughout the district. Also at Plaw
Hatch, Hatfield Forest, Manuden (Vicarage garden), and elsewhere.

Pinaceae

569. Pinus sylvestris L. : Scotch Fir In gardens; also near the Shell House in Hatfield Forest.

570. Pinus nigra Arnold : Corsican Pine In gardens.

571. Pinus wallichiana Jackson : BHUTAN PINE In Dunmow Road, Bishop's Stortford.

Page Fifty-five

- 572. Pinus pinea L. : STONE PINE A fine specimen in Elgin Coppice at Hatfield Forest.
- 573. Larix decidua Mill. : LARCH
 Single trees in gardens; occasional plantations. At Thorley;
 Hockerill Halt.
 - 574. Picea abies (L.) Karsten : Norway Spruce Common in plantations and gardens.
 - 575. Picea omorika Bolle : SERBIAN SPRUCE In gardens; Plaw Hatch.
 - 576. Abies alba Mill. : European Silver Fir In gardens and plantations.
- 577. Cedrus libani Loud. : CEDAR OF LEBANON
 Plaw Hatch; Hockerill Training College; Gilston Park; and in
 other grounds and gardens.
 - 578. Cedrus deodara Loud. : DEODAR In Elgin Coppice at Hatfield Forest; Hockerill Training College.
 - 579. Cedrus atlantica Manetti : ATLAS CEDAR In parks and gardens; Hatfield Forest.
 - 580. Cupressus macrocarpa Hart. : Monterey Cypress Common in gardens.
 - 581. Chamaecyparis lawsoniana (Murray) Parlatore : Lawson's Cypress Common in gardens.
 - 582. Sequoia wellingtonia Seeman : Mammoth Tree In gardens; frequent in this area.
 - 583. Sequoia sempervirens (Lambert) Endlicher : REDWOOD In gardens; frequent in this area.
- .584. Araucaria araucana (Molina) K. Koch : Monkey Puzzle Occasional in gardens, becoming rare.
- 585. Taxodium distichum Rich. : DECIDUOUS CYPRESS Elgin Coppice at Hatfield Forest; St. Mary's Convent, Bishop's Stortford.

Page Fifty-six

PTERIDOPHYTA

Ophioglossaceae

586. Ophioglossum vulgatum L. : Adder's Tongue Planting's Wood.

Polypodiaceae

- 587. Pteridium aquilinum (L.) Kuhn: BRACKEN
 There are colonies of it in various woods in our district but no
 extensive areas of it. Long Meadow; Brewery Fields; Bloodhounds
 Wood, etc.
 - 588. Dryopteris filix-mas agg. : MALE FERN In most of the woods in this district, sparingly.
 - 589. Dryopteris spinulosa (Mull.) Warr: NARROW PRICKLY-TOOTHED FERN
 Hatfield Forest (Spittlemore Coppice); Broome Wood.

Equisetaceae

- 590. Equisetum telmateia Ehrh. : Great Horse-Tail Hatfield Forest, at the end of the lake near Shell House.
 - 591. Equisetum arvense L. : Common Horse-Tail Frequent in waste places.
 - 592. Equisetum palustre L. : MARSH HORSE-TAIL Common in wet places. Hatfield Forest, in the swamp, etc.

Addendum

43a. Coronopus didymus (L.) Sm. : Lesser Wartcress (Swinecress)

A weed in a garden in Maple Avenue, 1951 (Ad).

- 55a. Saponaria officinalis L.: SOAPWORT
 Henham, in sand-pits by the roadside (Cd). Possibly not truly
 wild as the site was long ago occupied by cottages. These were pulled
 down twenty years ago and the plant has been there ever since. "Has
 colonized my garden in Windhill" (An).
 - 557a. Festuca ovina L. : Sheep's Fescue Hatfield Forest; Patmore Heath, 1951.

Page Fifty-seven